

Northeast Gulf Science

Volume 9
Number 1 *Number 1*

Article 6

4-1987

Occurrence of *Mysidopsis bahia* (Mysidacea: Mysidae) on the Atlantic Coast of Florida

Geraldine M. Cripe
U.S. Environmental Protection Agency

DOI: 10.18785/negs.0901.06

Follow this and additional works at: <https://aquila.usm.edu/goms>

Recommended Citation

Cripe, G. M. 1987. Occurrence of *Mysidopsis bahia* (Mysidacea: Mysidae) on the Atlantic Coast of Florida. *Northeast Gulf Science* 9 (1).
Retrieved from <https://aquila.usm.edu/goms/vol9/iss1/6>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

OCCURRENCE OF *Mysidopsis bahia* (MYSIDACEA: MYSIDAE) ON THE ATLANTIC COAST OF FLORIDA

Twenty-two specimens of *Mysidopsis bahia* (Molenock, 1969): 15 females, 4 males, and 2 juveniles (sex undetermined), were collected from Link Port Channel, Ft. Pierce, Florida on December 6, 1984, at 20‰ salinity and 24°C. Gravid females averaged 7 mm length (base of eyestalk to posterior ends of uropods, excluding setae) and carried an average of 5.4 young (range 2 to 10). Mature males ranged from 6 to 7 mm length (mean 6.5 mm). Dr. Thomas E. Bowman at the National Museum of Natural History, Smithsonian Institution, verified the identification.

Mysidopsis bahia was initially described from a collection taken from West Bay, Galveston, Texas (Molenock, 1969). It is distinguished from *M. almyra* and *M. bigelowi* by having 4 to 5 pairs of strong spines on the apex of the telson and 2 to 3 spines on the endopod of the uropod near the statocyst, (Molenock, 1969). To date, other specimens of this species have been taken from Laguna de Tamiahua, Veracruz, Mexico (Price, 1978), Cape Sable, Florida (Brattegard, 1970), and coastal Texas (Price, 1982), Louisiana, Mississippi and Alabama (Stuck *et al.*, 1979). To my knowledge, this is the first published report of this species from the Atlantic coast of Florida.

ACKNOWLEDGMENTS

I thank Dr. James R. Clark for collection and transportation of the specimens from the collection site, and Dr. T. Bowman for species' verification.

LITERATURE CITED

- Brattegard, T. 1970. Marine biological investigations in the Bahamas. 11. Mysidacea from shallow water in the Bahamas and southern Florida. Part 2. Sarsia 41:1-35.
- Molenock, J. 1969. *Mysidopsis bahia*, a new species of mysid (Crustacea: Mysidacea) from Galveston Bay, Texas. Tulane Stud. Zool. Bot. 15(3):113-116.
- Price, W.W. 1978. Occurrence of *Mysidopsis almyra* Bowman, *M. bahia* Molenock and *Bowmaniella brasiliensis* Bacescu (Crustacea: Mysidacea) from the eastern coast of Mexico. Gulf Res. Rept. 6(2):173-175.
- _____. 1982. Key to the shallow Mysidacea of the Texas coast with notes on their ecology. Hydrobiologia 93:9-21.
- Stuck, K.C., H.M. Perry and R.W. Heard. 1979. Records and range extensions of Mysidacea from coastal and shelf waters of the eastern Gulf of Mexico. Gulf Res. Rept. 6(3):239-248.

Environmental Research Laboratory/Gulf Breeze
Contribution Number 560.

Geraldine M. Cripe, U.S. Environmental Protection Agency, Environmental Research Laboratory, Sabine Island, Gulf Breeze, FL 32561.